

COVID-19 CLINICAL FEATURES OF THE PERIOD AND CHILL

Yuldasheva D. Yu.

Scientific Head, Assistant Professor, Tashkent Medical Academy Tashkent, Uzbekistan

Shaymuradova M.U

Master, Tashkent Medical Academy Tashkent, Uzbekistan

Abstract

The actual ministry. In the year of 2019 appeared which new koronavirus COVID-19 disease in the entire world across Sin the field of storage og'liqni serious problems causing it. SOVID19 the virus pregnancy and chill period, complicated night cause be can (Inoyatova a. sh. i dr. 2020.) The disease proliferation in the body at the time of pregnancy to stop or to defect, the mother of death increase and serious complications, the likelihood to come the reason is: the main disease severity and in the aftermath of them came out that the complications - respiratory failure, obstetrics blood is going to bea fetus intranatal death, the salamiis still the next purulent-septic diseases, the development lead cause can.

Keywords: - Agar COOLED-19 diagnosis confirmation - of the disease in a mild form with pregnant women therapist supervision under at home treatment extend, we can. Kaslightly of moderate and severe forms with you to the hospital and laid to required are.

Introduction:-

It's ay in the case, the Russian health ministry pregnant woman separated from to be employment recommend it. COOLED-19 knowan infected women, the majority , a few days after I passed away, who light a'lomatlarg a face will come. Ammo itself bad it feels too can. The results of research goals and objectives. Koronavirus infection COOLED-19 with infected pregnant in the chill of the period of clinical to the night the properties of the study. Materials and survey methods. 2021 is the year of Tashkent Medical Academy COVID-19 with the incidence of treatment towards

who came 40 students are pregnant. This pregnant 1-main group (1-AG) , as has been identified. Anamnezida the COVID-19 with kasalanmagan and Tashkent Medical Academy of pregnancy pathology of treatment for the purpose of who come pregnant 40 students, 2-control group (2-ng) as has been identified. Total research pregnant n=80 are. The survey results. Research 1-ag 40 students of pregnant aksaryati 34 students (85%) in term of the best choiceqlar if you pass, 6 students (15%) obstetrics guidelines with cut cutting operation was conducted. Obstetrics guidelines, in the uterus scar malfunction 3 students (50%) of the fetus uncertain quality 2 students (33,3%) and the pelvis, the head disproportsiyasi with 1 students (16,7%) to cut a cutting operation was conducted. Chill in the period , the best choiceis still the next 5 days complications and cut a cutting operation after 7 days in stasionar and chill period, the remaining days to live place check out to did not go. Of them, 3 students (6%) of the uterus subinvalyusiyasi observed, the reason that when they found out of the vagina 3 level of clean anamnezida which sexual organs of inflammatory diseases with kasallanganligini and often shamollab O'RVI with aching he'd said. 1 students (2%) of the salami of the 10-day body of xarorat that they had increased and the blood late to go to a complaint complications again make contact to come. The survey results according to the uterus subinvalyusiyasi and endometrid ULTRASOUND detected check of the uterus in the cavity of 25 mm, a blood clot will determine. This women of the uterus inside vakum make aspirasiya cleanses and uteratoniklar, antibiotics with treatment. Ultimately , the patient recover went to. A total of 4 persons (8%) gave birth in women who have uterine subinvalyusiyasi, 1 students (2%) body of xarorat increased and again 1 students (2%) late blood cutting were observed. Cut a cutting operation of 6 students (12%) of the women 2 students (33,3%) which operation after 1- and 2- of the day, the body of xarorat 38 °S to be up to were observed. In them the mainqa complications would be. Cut a cutting operation the all the women in protocol bo'yicha of antibiotics at the time of the operation only 2 g were sent to the meter 1 times with 100 ml of the vessel into drop many were sent to. Which operation after the body only xarorat of over 2 persons chill in the period to women antibiotics and send it to continue was held. 2-pregnant control group in the salamiq, 43 students (86%) in the period tug'ruqni is if you pass and 7 persons (14%) to cut a cutting operation with was completed. Cut a cutting operation instructions follows if. 4 students (57,1%) in the uterus scar malfunction, 1 students (14,3%) of the fetus, uncertain quality, 1 students (14,3%) of the pelvis disproportsiyasi and again 1 students (14,3%) preeklampsiyaning severe level (rightin the way of complications is ready is not) on the was held. Chill during the

period of childbirth women 1 students (2%) uterus subinvalyusiyasi observed, that women in the vagina with 3 level of clean and the anamnezida sex members inflammatory disease with the incidence competence for the treatment of known is. Homeqa complications were observed. Cut a cutting operation , then this control group 2 students (28,6%) women in the operation the next 1 - day body xarorat handle 38 °S to be increased were observed and in them antibiotics with treatment did not held. This is despite them next day the body of xarorat increase did not follow. So do 1-ag, chill period complications in the uterus subinvalyusiyasi 4 in (8%), cut a cutting operation , then the body of xarorat increased 2 students (33,3%) and tug'ruqdan then 1 students (2.3%), and chill in the period of late blood cutting 1 students (2%) was. 2-in ng, complications from next uterus subinvalyusiyasi 1 students (2%) and cut a cutting operation , then the body of xarorat increased 2 students (28,6%) were observed. Conclusions. COVID-19 , with the incidence past pregnant in the chill of the period of complications more it is, the uterus subinvalyusiyasi 8 %. The salamiis still the next purulent-septic diseases, the development of 6% came out. Chill in the period of late blood cutting to 2% up to make. Each both in the group cut a cutting operation for the reason of obstetrics guidelines are.

References

1. World Health Organization. Coronavirus disease (COVID-19) technical guidance: Infection prevention and control/WASH. Available at: <https://www.who.int/emergencies/diseases/novel-coronavirus-19/technical-guidance/infection-prevention-and-control>.
2. Временные методические рекомендации по профилактике, диагностике и лечению новой коронавирусной инфекции 2019-nCoV. Минздрав России. Версия 6 (24.04.2020). 142 с. Режим доступа: https://static-1.rosminzdrav.ru/system/attachmentsattaches/000/050/122/original/28042020_MR_COVID-19_v6.pdf. [Vremennye metodicheskie rekomendacii. "Profilaktika, diagnostika i lechenie novoj koronavirusnoj infekcii COVID-19", 6 versiya(28.04.2020). Available at: <http://nasci.ru/?id=11052>. Accessed: 1 june 2020. (In Russ).
3. Вступительное слово генерального директора ВОЗ на брифинге для средств массовой информации по КОВИД-19 - 11 марта 2020 года. <https://www.who.int/dg/speeches/detail/who-director-general-s-openingremarks-at-the-media-briefing-on-covid-19-11-march-2020>.

4. Ву Джей Ти, Леунг К. Леунг Г. М. Nowcasting and forecasting the potential domestic and international spread of the 2019-ncov outbreaking in Wuhan, China: A Modeling study. *Ланцет*. 2020; 395(10225): 689-697.
5. Гуань У. Ни З. Ху Й. et al. Клиническая характеристика коронавирусной болезни 2019 года в Китае. *N Engl J Med*. 2020; ([Epub впереди печати]).
6. Всемирная организация здравоохранения (ВОЗ) Доклад о ситуации с коронавирусной болезнью ВОЗ за 2019 год (COVID-19) 46. (Доступно по адресу:) https://www.who.int/docs/default-source/coronaviruse/situation-reports/20200306-sitrep-46-covid-19.pdf?sfvrsn=96b04adf_2 Дата обращения: 14 марта 2020 года.
7. Нельсон - Пирси К. Респираторное заболевание. в: *Справочник по акушерской медицине*. CRC Press, Бока-Ратон, Флорида 2015: 63-84.
8. Гарднер М.О., Дойл Н. М. Астма во время беременности. *Акушер-Гинеколог Клин Норт Ам*. 2004; 31: 385-413.
9. Ши Х. Хан Х. Цзян Н. et al. Рентгенологические данные 81 пациента с пневмонией COVID-19 в Ухане, Китай: описательное исследование. *Ланцет Дис*. 2020; 20: 425-434.
10. Wong C.K., Lam CWK, Wu AKL, Ip WK, Lee NL, Chan IH и др. Плазменные воспалительные цитокины и хемокины при тяжелом остром респираторном синдроме (ОРВИ) *Clin Exp Immunol*. 2004;136(1):95–103.[PMC free article] [PubMed] [Google Scholar].